



From Nothing to the Universe

V. Mukhanov

LMU, München

The efforts to understand the universe is one of the very few things that lifts human life a little above the level of farce...

S. Weinberg, 1977



$$\Delta q \times \Delta p \geq \frac{1}{2} \hbar$$

Before 1990

Notre concitoyen, disaient-ils en pleurant,
Perd l'esprit : la lecture a gâté Démocrite.
Nous l'estimerions plus s'il était ignorant.
Aucun nombre, dit-il, les mondes ne limite :
 Peut-être même ils sont remplis
 De Démocrites infinis.

La Fontaine

“Our fellow citizen,” they said, “has lost his mind”
Reading has ruined Democritus.
If he knew less he’d have more sympathy from us.
There are more worlds, he claims, in number infinite,
 And each of them may have in it
 Another Democritus.


La Fontaine

“Only by their breaking could the divine configurations be perfected”

Kabbalistic text; Ta'alumoth Chokhmah (The Channels of Wisdom)
1629, Joseph Samomon del Medigo of Crete













● The Universe expands



● Hubble law



$$v = Hr$$

$$t \sim \frac{r}{v} = \frac{1}{H} \sim 13,7 \text{ bil. years}$$



There is baryonic matter:


about 25% of ^4He , D...heavy elements

Dark Matter???? baryonic origin???



Large Scale Structure: clusters of galaxies!


Filaments, Voids????????????????????????????????

- There exists background radiation with the temperature $T \approx 3K$




Penzias, Wilson 1965





a





$$\lambda \propto a$$




$$T \propto \frac{1}{a}$$


When the Universe was 1000 times smaller
its temperature was about $2725^{\circ}K$

Big Bang




Big Bang






Millennium Simulation

10,077,696,000 particles




Big Bang



Very homogeneous

Inhomogeneous



$$\frac{\text{female}}{\text{male}} = 0.96$$

Quantum fluctuations and a nonsingular Universe

V.F. Mukhanov and G.V. Chibisov

P. N. Lebedev Physics Institute, Academy of sciences of the USSR

(Submitted 26 February 1981; 15 April 1981)

Adopting a perturbation of the curvature scalar as a physical variable, we find the corresponding action in the form [6]

$$\delta S_b = \frac{1}{2} \int d^4x \left[\dot{\phi}^2 - \nabla^\alpha \phi \nabla_\alpha \phi + \left(\frac{a''}{a} + M^2 a^2 \right) \phi^2 \right], \quad (5)$$

where $\phi = 1/\sqrt{18(4H^2 - M^2)} a \delta R / M \ell$, and $\ell = (8\pi G/3)^{1/2} = 4.37 \times 10^{-33}$ cm is the Planck length.


A finite duration of the de Sitter stage does not by itself rule out the possibility that this stage may exist as an intermediate stage in the evolution of the universe. An interesting question arises here: Might not perturbations of the metric, which would be sufficient for the formation of galaxies and galactic clusters, arise in this stage? To answer this question, we need to calculate the correlation function for the fluctuations of the metric after the universe goes from the de Sitter stage to the hydrodynamic stage. By analogy with (6) we find

$$\langle 0 | \hat{h}(\mathbf{x}) \hat{h}(\mathbf{x} + \mathbf{r}) | 0 \rangle = \frac{1}{2\pi^2} \int Q^2(k) \frac{\sin kr}{kr} \frac{dk}{k}, \quad (8)$$


where $h = h_\alpha^\alpha$ and where, for the most interesting region, $H > k > H \exp(-3H^2/M^2)$ ($M^2 \ll H^2$),

$$Q(k) \approx 3\ell M \left(1 + \frac{1}{2} \ln \frac{H}{k} \right). \quad (9)$$



The fluctuation spectrum is thus nearly flat. The quantity $Q(k)$ is the measure of the amplitude of perturbations with scale dimensions $1/k$ at the time the universe begins the ordinary Friedmann expansion. With $\ell M \sim 10^{-3} - 10^{-5}$ and $M/H \leq 0.1$ —these values are consistent with modern theories of elementary particles—the amplitude of the perturbations of the metric on the



$n_s = 0.96$



Very homogeneous



Inhomogeneous





$$\Delta x m \Delta v \geq \hbar$$



$$\rightarrow \Delta p \Delta x \geq h$$

↓

There always exist **unavoidable**
Quantum Fluctuations




Quantum fluctuations in the density distribution are large (10^{-5})

only in extremely small scales ($\sim 10^{-33}$ cm),

but very small ($\sim 10^{-58}$) on galactic scales ($\sim 10^{25}$ cm)

Can we transfer the large fluctuations from extremely small scales to large scales???



The space and time had both one beginning . The space was made not in time but simultaneously with time.

Saint Augustin of Hipo

God creates new worlds constantly

Zohar

If the world has begun with a single quantum, the notions of space and time would altogether fail to have any sense at the beginning and would only begin to get some sensible meaning when the original quantum would have been divided in a sufficient number of quanta.

G. Lemaître,

The beginning of the world from the point of view of quantum theory, Nature 127, 706 (1931)


ANNALS OF PHYSICS 115, 78–106 (1978)

The Creation of the Universe as a Quantum Phenomenon

R. BROUT, F. ENGLERT, AND F. GUNZIG

Faculté des Sciences, Université Libre de Bruxelles, Bruxelles, Belgium

Received July 7, 1977




Predictions!!!

1)

Does space have a shape?

LD © 2008 HowStuffWorks

Euclidian
Space




Zero Curvature

Elliptical
Space



Positive Curvature

Hyperbolic
Space




Negative Curvature


$$\Omega = 1$$

Perturbations (inhomogeneities) are:

2) Adiabatic (MC 1981)



100 photons
50 baryons



98 photons
49 baryons


$$~~49 - 2 = 47~~$$

3) Gaussian (MC 1981)



$$\Phi = \Phi_g + f_{NL}\Phi_g^2, \text{ where } f_{NL} = O(1) \text{ (MC, 81)}$$

4) have log spectrum (MC 1981)



I I I

Amplitude increases by a few percents when scale increases in few times

$$4) \Phi \propto \ln(\lambda/\lambda_\gamma) \propto \lambda^{1-n_s} \text{ with } n_s = 0.96 \text{ (MC, 1981)}$$

L.P. 9/6/2003:

We are writing a proposal to get money to do our small angular scale CMB experiment. If I say that simple models of inflation require $n_s=0.95\pm 0.03$ (95% cl) is it correct?

I'm especially interested in the error. **Specifically, if $n_s=0.99$ would you throw in the towel on inflation?**

V.M. 9/8/2003






The "robust" estimate for spectral index for inflation is $0.92 < n_s < 0.97$.

The upper bound is more robust than lower. The physical reason for the deviation of spectrum from the flat one is the necessity to finish inflation....




If you find $n_s=0.99 \pm 0.01$ (3 sigma) I would throw in the towel on inflation.


After 90 - present




COBE 1992



Local Experiments as of 1999
[calibration error not included]







1965




Penzias and Wilson




1992




COBE



2003



WMAP




2009



Planck

???
End 2012



the Planck Collaboration, including individuals from more than 100 scientific institutes in Europe, the USA and Canada



planck



DTU Space
National Space Institute



Science & Technology
Facilities Council



CSIC



Hfi PLANCK
to look back to the birth of the Universe



National Research Council of Italy






DLR Deutsches Zentrum
für Luft- und Raumfahrt e.V.



Planck is a project of the European Space Agency, with instruments provided by two scientific Consortia funded by ESA member states (in particular the lead countries: France and Italy) with contributions from NASA (USA), and telescope reflectors provided in a collaboration between ESA and a scientific Consortium led and funded by Denmark.



WMAP





Planck





THUW
SCI





Cobe 1992

WMAP 2003

Planck 2013

PREDICTIONS

1) flat Universe


Perturbations are :


2) adiabatic (MC, 81)

3) gaussian: $\Phi = \Phi_g + f_{NL} \Phi_g^2$, where $f_{NL} = \mathcal{O}(1)$ (MC, 81)


4) spectrum: $\Phi \propto \ln(\lambda/\lambda_\gamma) \propto \lambda^{1-n_s}$ with $n_s = 0.96$ (MC, 81)


with $\Omega_{tot} = 1$ (prediction) and H_0 , Ω_{Λ} , Ω_{bar} from supernova, deuterium et.cet. we get








Angular scale











Before Planck



After Planck

A Cosmic Microwave Background (CMB) fluctuation map, showing temperature variations across the sky. The map is an oval shape with a complex, noisy pattern of blue and orange colors, representing different temperature fluctuations. The background is a dark field of stars.

$-\Omega_{tot} = 1 \pm 0.005$




– Perturbations are adiabatic

– Gaussian: $f_{NL} = 2 \pm 5$

$n_s = 0.96 \pm 0.005!!!$

CONCLUSIONS

- General Relativity is valid up to the scales 10^{-27} cm
- We all originated from quantum fluctuations



- Theory is right
- Planck is right
- BICEP2 is right

$T+P \quad \checkmark \quad T+B \quad \checkmark$

$P+B \quad \checkmark$

but

~~$T+P+B$~~

Therefore $P+B \Rightarrow$ catastrophe for
theory



29. April 2014 10:54 Entstehung des Universums

Risse in der Urknall-Theorie



Forschungsstation am Südpol: Hier meinen Physiker Signale aus den ersten Sekundenbruchteilen nach dem Urknall gemessen zu haben. Viele Kollegen sind noch nicht überzeugt. (Foto: REUTERS)

Signale aus der Geburtsstunde des Universums: Mitte März jubelte ein Forscherteam über eine bahnbrechende Messung von Gravitationswellen. Möglicherweise haben die Physiker sich zu früh gefreut.

Von *Marlene Weiß*

Diskutieren
Versenden
Drucken
Wer meint, die Welt erklären zu können, indem er am kleinen π schraubt, bekommt es mit Viatcheslav Mukhanov zu tun. "Vollkommener Unsinn", schimpft der an der Uni München aktive russische Physiker, "die Zeitschriften sind voll davon, aber es bleibt trotzdem Unsinn!"



Feedback

Auch wer sonst nichts von seinem Vortrag kürzlich am Max-Planck-Institut für Astrophysik in Garching bei München verstanden hat, eines dürfte jedem Zuhörer klar geworden sein: Das kleine π in den Formeln über den Beginn des Universums, auch "spektraler Index" genannt, sollte man in Ruhe lassen, wenn man sich nicht mit Mukhanov anlegen möchte.

Das sind schlechte Nachrichten für all die Fachleute, die Mitte März jubelten, als es hieß, man habe mit einem Teleskop am Südpol Signale aus den ersten Sekundenbruchteilen nach dem Urknall gemessen: Vielleicht war der Jubel verfrüht, das Ergebnis widerspricht anderen Messungen.

Spuren von Gravitationswellen, die vor 13,82 Milliarden Jahren entstanden sein

ANZEIGE

Popmusik aus Bayern und dem Alpenraum amazon.de

